



注意事項：

- 請列出計算過程。否則，不予計分。
- 如需要，可用所附表格求值。
- 請務必將題號標示清楚。可不按順序作答。

1. (15%) The manager of a certain pharmacy claims that the times to process orders are exponentially distributed with a mean of 12 minutes. If 100 customers visit the pharmacy in a 2-day period, and if there are 50, out of these 100, customers waiting for more than 12 minutes, what is your comment on the manager's claim? Give your probabilistic reason.
2. (20%) Let Y_1, \dots, Y_n be a random sample of size n from a population which is uniformly distributed over (θ_1, θ_2) with unknown θ_1 and θ_2 . Also let $Y_{(1)} = \min(Y_1, \dots, Y_n)$ and $Y_{(n)} = \max(Y_1, \dots, Y_n)$
 - (a) Is $Y_{(1)}$ a consistent estimator for θ_1 ? Why?
 - (b) Is $Y_{(n)}$ a consistent estimator for θ_2 ? Why?
 - (c) Show that $Y_{(1)}$ and $Y_{(n)}$ are jointly sufficient for θ_1 and θ_2 .
3. (20%) Let Y_1, \dots, Y_n be a random sample from a population with density function given by

$$f(y) = \begin{cases} \left(\frac{1}{\theta}\right) y^{\theta-1} e^{-y/\theta}, & y > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Find a uniformly most powerful test for testing $H_0: \theta = \theta_0$ against $H_a: \theta > \theta_0$, with significance level α .
- (b) Suppose that 5 random observations from this population were taken as follows.

26.6 31.3 27.3 33.4 29.7

Use the result of part(a) to test $H_0: \theta = 10$ versus $H_a: \theta > 10$, with $\alpha = 0.05$.

- (c) Estimate the p -value of the observed data in part(b) and explain it.
4. (20%) A survey was conducted to compare the fraction of students favoring candidate A in four universities. Random samples of 200 students were polled in each of the four universities, with the results as shown in the following table.

Opinion	University				Total
	1	2	3	4	
Favor A	77	52	61	46	236
Do not favor A	123	148	139	154	564
Total	200	200	200	200	800

Suppose that the numbers of students favoring candidate A in the four samples can be regarded as four independent binomial random variables. Construct a likelihood ratio test, with significance level $\alpha = 0.05$, of the hypothesis that the fractions of students favoring candidate A are the same in all four universities.

5. (25%) An experiment was conducted to determine the effect of input x on the output y . One run of the experiment at each input setting gave the following data.


x	-3	-2	-1	0	1	2	3
y	1	0	0	-1	-1	0	0

- (a) Fit the model $Y = \beta_0 + \beta_1 x + \beta_2 x^2 + \epsilon$.
- (b) Does the quadratic input significantly affect the output? Test this with significance level $\alpha = 0.05$. Also estimate the p -value and explain it.
- (c) The experimenter claims that the minimum value of $E(Y)$ occurs at $x = 1$. Test this claim at significance level $\alpha = 0.05$.
- (d) Find a 95% confidence interval for the mean output at $x = 0$.

參考用

Table 1

Normal curve areas
Standard normal probability in right-hand
tail for negative values of z areas are found
by symmetry



Second decimal place of z

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0515	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
2.8	.0026	.0025	.0024	.0023	.0022	.0022	.0021	.0021	.0020	.0019
2.9	.0019	.0018	.0017	.0017	.0016	.0016	.0015	.0015	.0014	.0014
3.0	.0013									
3.5	.000233									
4.0	.0000317									
4.5	.00000340									
5.0	.000000287									

From R. E. Walpole, *Introduction to Statistics* (New York: Macmillan, 1963)

Table 2
Percentage points of the t-distribution



t _{0.95}	t _{0.90}	t _{0.85}	t _{0.80}	t _{0.75}	t _{0.70}	t _{0.65}	t _{0.60}	t _{0.55}	t _{0.50}	df.
3.078	6.314	12.706	31.821	63.657	1					
1.886	2.920	4.303	6.965	9.925	2					
1.638	2.353	3.182	4.541	5.841	3					
1.533	2.132	2.776	3.747	4.604	4					
1.476	2.015	2.571	3.365	4.032	5					
1.440	1.943	2.447	3.143	3.707	6					
1.415	1.895	2.365	2.998	3.499	7					
1.397	1.860	2.306	2.896	3.355	8					
1.383	1.833	2.262	2.821	3.250	9					
1.372	1.812	2.238	2.764	3.169	10					
1.363	1.796	2.201	2.718	3.106	11					
1.356	1.782	2.179	2.681	3.055	12					
1.350	1.771	2.160	2.650	3.012	13					
1.345	1.761	2.145	2.624	2.977	14					
1.341	1.753	2.131	2.602	2.947	15					
1.337	1.746	2.120	2.583	2.921	16					
1.333	1.740	2.110	2.567	2.898	17					
1.330	1.734	2.101	2.552	2.878	18					
1.328	1.729	2.093	2.539	2.861	19					
1.325	1.725	2.086	2.528	2.845	20					
1.323	1.721	2.080	2.518	2.831	21					
1.321	1.717	2.074	2.508	2.819	22					
1.319	1.714	2.069	2.500	2.807	23					
1.318	1.711	2.064	2.492	2.797	24					
1.316	1.708	2.060	2.485	2.787	25					
1.315	1.706	2.056	2.479	2.779	26					
1.314	1.703	2.052	2.473	2.771	27					
1.313	1.701	2.048	2.467	2.763	28					
1.311	1.699	2.045	2.462	2.756	29					
1.282	1.645	1.960	2.326	2.576	inf.					


From "Table of Percentage Points of the t-Distribution,"
Computed by Maxine Merrington, *Biometrika*, Vol. 33 (1941), p.
309. Reproduced by permission of Professor E. S. Pearson.

表 1
表 2

參考用

Table 3

Percentage points of the χ^2 distributions



d.f.	$\chi^2_{0.995}$	$\chi^2_{0.990}$	$\chi^2_{0.975}$	$\chi^2_{0.950}$	$\chi^2_{0.900}$
1	0.0000393	0.0001571	0.0009821	0.0039321	0.0157908
2	0.0100231	0.0201007	0.0506356	0.102387	0.210720
3	0.0717212	0.114832	0.215795	0.351846	0.584375
4	0.206990	0.297110	0.484419	0.710721	1.065623
5	0.411740	0.554300	0.831211	1.145476	1.61031
6	0.675727	0.872085	1.237347	1.63539	2.20413
7	0.989265	1.239043	1.68987	2.16735	2.83311
8	1.344419	1.646482	2.17973	2.73264	3.48954
9	1.734926	2.087912	2.70039	3.32511	4.16816
10	2.15585	2.55821	3.24697	3.94030	4.86518
11	2.60331	3.05347	3.81575	4.57481	5.57779
12	3.07382	3.57056	4.40379	5.22603	6.30380
13	3.56503	4.10691	5.00874	5.89186	7.04150
14	4.07468	4.66043	5.62872	6.57063	7.78953
15	4.60094	5.22935	6.26214	7.26094	8.54675
16	5.14224	5.81221	6.90766	7.96164	9.31223
17	5.69724	6.40776	7.56418	8.67176	10.0852
18	6.26481	7.01491	8.23075	9.39046	10.8649
19	6.84398	7.63273	8.90655	10.1170	11.6509
20	7.43386	8.26040	9.59083	10.8508	12.4476
21	8.03366	8.89720	10.28293	11.5913	13.2536
22	8.64272	9.54249	10.9823	12.3380	14.0715
23	9.26042	10.19567	11.6885	13.0905	14.8979
24	9.88623	10.8564	12.4011	13.8484	15.6587
25	10.5197	11.5240	13.1197	14.6114	16.4734
26	11.1603	12.1981	13.8439	15.3791	17.2919
27	11.8076	12.8786	14.5733	16.1513	18.1138
28	12.4613	13.5648	15.3079	16.9279	18.9392
29	13.1211	14.2565	16.0471	17.7083	19.7677
30	13.7867	14.9535	16.7908	18.4926	20.5992
40	20.7065	22.1643	24.4331	26.5093	29.0505
50	27.9907	29.7067	32.3574	34.7642	37.6886
60	35.5346	37.1848	40.4817	43.1879	46.4589
70	43.2752	45.4418	48.7576	51.7293	55.3290
80	51.1720	53.5400	57.1532	60.3915	64.2778
90	59.1963	61.7541	65.6466	69.1260	73.2912
100	67.3276	70.0648	74.2219	77.9295	82.3581

頁三第

Table 3 (continued)

d.f.	$\chi^2_{0.100}$	$\chi^2_{0.075}$	$\chi^2_{0.050}$	$\chi^2_{0.025}$	$\chi^2_{0.010}$	$\chi^2_{0.005}$
1	2.70554	3.84146	5.02389	6.63490	7.87944	
2	4.60517	5.99147	7.37776	9.21034	10.5966	
3	6.25139	7.81473	9.34840	11.3449	12.8381	
4	7.77944	9.48773	11.1433	13.2767	14.8602	
5	9.23635	11.0705	12.8325	15.0863	16.7496	
6	10.6446	12.5916	14.4494	16.8119	18.5476	
7	12.0170	14.0671	16.0128	18.4753	20.2777	
8	13.3616	15.5073	17.5346	20.0902	21.9550	
9	14.6837	16.9190	19.0228	21.6660	23.5893	
10	15.9871	18.3070	20.4831	23.2093	25.1882	
11	17.2750	19.6751	21.9200	24.7250	26.7569	
12	18.5494	21.0261	23.3367	26.2170	28.3095	
13	19.8119	22.3621	24.7356	27.6883	29.8194	
14	21.0642	23.6848	26.1190	29.1413	31.3193	
15	22.3072	24.9958	27.4884	30.5779	32.8013	
16	23.5418	26.2962	28.8484	31.9999	34.2672	
17	24.7690	27.5871	30.1910	33.4087	35.7185	
18	25.9884	28.8693	31.5264	34.8053	37.1564	
19	27.2036	30.1435	32.8523	36.1908	38.5822	
20	28.4120	31.4101	34.1696	37.5662	39.9968	
21	29.6151	32.6705	35.4789	38.9331	41.4010	
22	30.8133	33.9244	36.7807	40.2894	42.7956	
23	32.0069	35.1725	38.0757	41.6384	44.1813	
24	33.1963	36.4151	39.3641	42.9798	45.5585	
25	34.3816	37.6525	40.6465	44.3141	46.9278	
26	35.5631	38.8852	41.9252	45.6417	48.2899	
27	36.7412	40.1133	43.1944	46.9650	49.6449	
28	37.9159	41.3372	44.4607	48.2782	50.9933	
29	39.0875	42.5569	45.7222	49.5879	52.3356	
30	40.2560	43.7729	46.9792	50.8922	53.6720	
40	51.8050	55.7585	59.3417	63.6907	66.7659	
50	63.1671	67.5048	71.4202	76.1539	79.4900	
60	74.3970	79.0819	83.2976	88.3794	91.9517	
70	85.5271	90.5312	95.0231	100.425	104.215	
80	96.5782	101.879	106.629	112.329	116.321	
90	107.565	113.145	118.136	124.116	128.299	
100	118.498	124.342	129.561	135.807	140.169	

From "Tables of the Percentage Points of the χ^2 -Distribution," *Biometrika*, Vol. 32 (1941), pp. 138-189, by Catherine M. Thompson. Reproduced by permission of Professor E. S. Pearson.