

## 國立中央大學104學年度碩士班考試入學試題

所別:<u>光電科學與工程學系碩士班不分組(一般生)</u>科目:近代物理 共 乙 頁 第 1 頁 本科考試可使用計算器,廠牌、功能不拘 \*請在答案卷(卡)內作答

- 1. What is the minimum photon energy required to create an e<sup>-</sup>e<sup>+</sup> pair when a photon collides with a free electron at rest? (10 pts)
- 2. If the protons were accelerated to a kinetic energy of 6.5 GeV, what magnetic field was needed to keep the protons traveling in a circle of 15 m? (10 pts)
- 3. A photon having 50 keV scatters from a free electron at rest. What is the maximum energy that the electron can obtain? (10 pts)
- 4. A muonic atom consists of a muon (mass m=106 MeV/c<sup>2</sup> and charge q=-e) in place of an electron. For the muon in a hydrogen atom, what is the binding energy of the muon in the ground state? (10 pts)
- 5. For waves with the phase velocity equal to the group velocity, derive the dependence of the phase velocity on the wavelength. (10 pts)
- 6. A quantum dot (QD) is a nanocrystal made of semiconductor materials that are small enough to exhibit quantum mechanical properties. Assuming a cubic quantum dot has length a on each side as shown in Fig.1. Please try to answer the following questions with clear explanations. A simple answer without explanation will not be scored.
  - a. (5 pts) Please use uncertainty principle to estimate the lowest possible energy of a free electron confined in this quantum dot.
  - b. (15 pts) Please use Schrödinger equation to obtain the general form of the wave functions and corresponding energy of the energy levels of a free electron confined in this quantum dot.
  - c. (5 pts) Explain why the above results of the lowest possible energy are identical or different.
  - d. (5 pts) If we applied an magnetic field B along one axis in such a quantum dot with considering the spin of electron(s=1/2,  $\mu_B = \frac{e\hbar}{2m_e}$ ), what is the energy difference between the two lowest possible electron energy levels?
- e. (5 pts) Is it possible for a electron has transition from the lowest possible energy level to the third lowest possible energy level by absorbing a photon?





## 國立中央大學104學年度碩士班考試入學試題

所別:光電科學與工程學系碩士班 不分組(一般生) 科目:近代物理 共 2 頁 第 2 頁 本科考試可使用計算器,廠牌、功能不拘 \*請在答案卷(卡)內作答




Fig. 1 A cubic quantum dot has length a on each side. Electron can only exist within this quantum dot.

- 7. (5 pts) In how many different ways can three identical particles be assigned to five states? Please consider both boson and fermion cases.
- 8. (5 pts) Does energy bands in solid follow selection rules? Why?
- 9. (5 pts) Give an example in solid for each bonding mechanism. (Van der Waals, hydrogen bond, covalent bond, and metallic bond)



注:背面有試題