類組:電機類 科目:工程數學 C(3005)

共12頁第一頁

## ※請在答案卡內作答

- · 本測驗試題為多選題 (答案可能有一個或多個) ,請選出所有正確或最適當的答案,並請用2B鉛筆作答於答案卡。
- 共二十題,每題五分。每題ABCDE每一選項單獨計分;每一選項的個別分數為一分,答錯倒扣一分。

Notation: In the following questions, underlined letters such as  $\underline{a}, \underline{b}$ , etc. denote column vectors of proper length; boldface letters such as A, B, etc. denote matrices of proper size;  $A^{\mathsf{T}}$  means the transpose of matrix A.  $I_n$  is the  $(n \times n)$  identity matrix.  $\|\underline{a}\|$  means the Euclidean norm of vector  $\underline{a}$ .  $\mathbb{R}$  is the usual set of all real numbers;  $\mathbb{C}$  is the usual set of all complex numbers. By  $A \in \mathbb{R}^{m \times n}$  we mean A is an  $m \times n$  real-valued matrix. u(x) is unit-step function defined as u(x) = 1 if  $x \geq 0$  and u(x) = 0 if x < 0;  $\star$  is the convolution operator;  $\mathcal{L}: f(x) \mapsto F(s)$  and  $\mathcal{L}^{-1}: F(s) \mapsto f(x)$  denote the unilateral Laplace and inverse Laplace transforms for  $x \geq 0$ , respectively.

- -- Let  $\mathbb{H} = \mathbb{I}_n 2\underline{u}\,\underline{u}^{\top}$ , where  $\underline{u} \in \mathbb{R}^n$ ,  $n \geq 2$  and  $\|\underline{u}\| = 1$ . Which of the following statements is/are true?
  - (A) The matrix H is both symmetric and orthogonal.
  - (B) Both 1 and -1 are eigenvalues of H.
  - (C) det(H) = 1.
  - (D) Trace(H) = n-2.
  - (E) None of the above.



# 類組:電機類 科目:工程數學 C(3005)

共12頁第2頁

#### ※請在答案卡內作答

- Two square matrices A and B are similar, denoted by  $A \sim B$ , if  $B = P^{-1}AP$  for some nonsingular matrix P. Which of the following statements is/are true?
  - (A) Two similar matrices always have the same set of eigenvalues, including multiplicity.
  - (B) Two  $n \times n$  matrices having the same set of eigenvalues, including multiplicity, are similar.
  - (C) Any two square matrices with the same trace and determinant are similar.
  - (D) If  $A \sim B$ , then  $p(A) \sim p(B)$  for any polynomial p(x).
  - (E) None of the above.

- $\Xi$ . Let  $\mathbf{A} = \underline{x} \underline{y}^{\mathsf{T}}$ , where  $\underline{x}$  and  $\underline{y}$  are two nonzero vectors of  $\mathbb{R}^n$ , n > 1. Which of the following statements is/are true?
  - (A) rank(A) = 1 and the range space of A is  $Span\{\underline{y}\}$ .
  - (B)  $\operatorname{nullity}(\mathbf{A}) = 2$  and the  $\operatorname{null} \operatorname{space} \operatorname{of} \mathbf{A} \operatorname{is} \operatorname{Span}\{\underline{x},\underline{y}\}.$
  - (C) Trace(A) = 1 and det(A) = 0
  - (D) A is always diagonalizable.
  - (E) None of the above.



注·背面有試題

類組:電機類 科目:工程數學 C(3005)

共12頁第2頁

※請在答案卡內作答

- $\underline{w}$ . Let  $\underline{A} = \underline{x}\underline{y}^{\mathsf{T}} + \underline{y}\underline{x}^{\mathsf{T}}$ , where  $\underline{x}$  and  $\underline{y}$  are two nonzero orthonormal vectors of  $\mathbb{R}^n$  and n > 2. Which of the following statements is/are true?
  - (A) Both  $\underline{x}$  and y are eigenvectors of A.
  - (B) Trace(A) = 1 and det(A) = 0.
  - (C) A is not diagonalizable.
  - (D) The least square solution of  $\mathbf{A}\underline{z} = \underline{b}$ , where  $\underline{b}$  is a vector in  $\mathbb{R}^n$ , is  $(\underline{b}^{\top}\underline{x})\underline{x} + (\underline{b}^{\top}\underline{y})\underline{y}$ .
  - (E) None of the above.



- £. Let  $A, B \in \mathbb{R}^{n \times n}$ , and  $\{\underline{u}_1, \dots, \underline{u}_n\}$  be an orthonormal basis for  $\mathbb{R}^n$ . It is known that  $\langle A, B \rangle = \operatorname{Trace}(A^{\top}B)$  is an inner product. We denote  $A \perp B$  if  $\langle A, B \rangle = 0$ . Which of the following statements is/are true?
  - (A) Let  $\underline{x}, \underline{y}, \underline{w}, \underline{z}$  be four nonzero vectors of  $\mathbb{R}^n$ . Then  $\underline{w} \underline{z}^{\top} \perp \underline{x} \underline{y}^{\top}$  if and only if  $\underline{w} \perp \underline{x}$  and  $\underline{z} \perp y$ .
  - (B) The set  $\mathcal{B}_1 := \left\{ \underline{u}_i \, \underline{u}_j^\mathsf{T} : i, j = 1, \dots, n \right\}$  is an orthonormal basis for  $\mathbb{R}^{n \times n}$ .
  - (C) The set

$$\mathcal{B}_2 = \left\{ \underline{u}_i \, \underline{u}_i^\top + \frac{\underline{u}_i \, \underline{u}_j^\top + \underline{u}_j \, \underline{u}_i^\top}{\sqrt{2}} : 1 \le i < j \le n \right\}$$

is an orthonormal basis for the real vector space  $\mathcal{S}_1 = \left\{ \mathbf{A} \in \mathbb{R}^{n \times n} : \mathbf{A} = \mathbf{A}^\top \right\}$ .

(D) The set

$$\mathcal{B}_3 = \left\{ \frac{\underline{u}_i \, \underline{u}_j^\top - \underline{u}_j \, \underline{u}_i^\top}{\sqrt{2}} : 1 \le i < j \le n \right\}$$

is an orthonormal basis for the real vector space  $\mathcal{S}_2 = \left\{ \mathbf{A} \in \mathbb{R}^{n \times n} : \mathbf{A} = -\mathbf{A}^\top \right\}$ .

(E) None of the above.

類組:電機類 科目:工程數學 C(3005)

共12頁第止頁

※請在答案卡內作答

 $\dot{\pi}$ 、 The system of linear equations  $\mathbf{A}\underline{x} = \underline{b}$  has

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}, \quad \underline{b} = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$$

Which of the following statements is/are true?

- (A) rank(A) + nullity(A) = 3.
- (B)  $\mathbf{A}^{\mathsf{T}}\mathbf{A}$  is a symmetric  $2 \times 2$  matrix.
- (C) The nullspace of A has two linearly independent vectors.
- (D)  $\mathbf{A}^{\mathsf{T}}\mathbf{A}$  is an invertible matrix.
- (E) None of the above.

七、 Continued from Problem 六, which of the following statements is/are true?

(A) The matrix A has the same row space as the following matrix

$$\left[\begin{array}{ccc} 1 & 1 & 1 \\ 0 & 1 & 2 \end{array}\right]$$

- (B)  $\det(\mathbf{A}\mathbf{A}^{\mathsf{T}}) = 0$ .
- (C) Let  $\underline{p}$  be the projected vector of  $\underline{b}$  onto the column space of  $\mathbf{A}$ . The Euclidean distance between  $\underline{b}$  and  $\underline{p}$  is zero.
- (D) There exists a  $(3 \times 2)$  matrix C such that rank(CA) = 3.
- (E) None of the above.



類組:電機類 科目:工程數學 C(3005)

共12頁第5頁

※請在答案卡內作答

 $\wedge$  Continued from Problem  $\Rightarrow$ , let  $B_1$  be a  $(2 \times 2)$  matrix and consider the system  $B_1A\underline{x} = B_1\underline{b}$ . It is known that

$$\mathbb{B}_1 \underline{b} = \begin{bmatrix} 5 \\ 0 \end{bmatrix} \text{ and } \underline{x} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

Which of the following vectors can be column vectors of  $B_1$ ?

- $(A) \begin{bmatrix} 1 \\ 0 \end{bmatrix}$
- (B)  $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$
- (C)  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$
- (D)  $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- (E) None of the above.



九、 Given the matrices

$$\mathbf{M}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad \mathbf{M}_2 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{M}_3 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

which of the following statements is/are true?

- (A)  $M_1$ ,  $M_2$  and  $M_3$  are linearly independent over  $\mathbb R$  in  $\mathbb R^{2\times 2}$ .
- (B) The span of  $\{M_1, M_2, M_3\}$  is the set of all  $(2 \times 2)$  real matrices.
- (C) The set of all Hermitian  $(2 \times 2)$  complex-valued matrices is a subspace of the span of  $\{M_1, M_2, M_3\}$  over  $\mathbb{C}$ .
- (D) Any linear combination of  $M_1$ ,  $M_2$  and  $M_3$  can be diagonalized over  $\mathbb C$ .
- (E) None of the above.



類組:<u>電機類</u> 科目:<u>工程數學 C(3005)</u>

共して真第しり真

※請在答案卡內作答

+ Continued from Problem  $\pi$ , let  $B=3M_1+4M_2+M_3$  and  $C=B^4-4B^3-9B^2+27B+111_2$ . Which of the following is/are true?

(A) 
$$C = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

(B) 
$$C = \begin{bmatrix} 3 & 0 \\ 0 & 4 \end{bmatrix}$$

(C) 
$$C = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

(D) 
$$C = \begin{bmatrix} 3 & 1 \\ 1 & 4 \end{bmatrix}$$

(E) None of the above.



 $+-\cdot$  Solve the first-order differential equation  $x^2y'(x)+xy(x)\ln(y(x))=xy(x)$ . Which of the following statements is/are true?

- (A) This is a homogeneous and linear differential equation.
- (B) y(x) = 0 is one particular solution.
- (C)  $y(x) = \exp(1)$  is another particular solution.
- (D) x = 0 is also a solution.
- (E) None of the above.

類組: <u>電機類</u> 科目: <u>工程數學 C(3005)</u>

共一つ頁第八頁

#### ※請在答案卡內作答

- $+\pm$ . Continued from Problem +-. Given the initial condition y(a)=b, which of the following statements is/are true?
  - (A) No solution if a = 0.
  - (B) A unique solution if b = a > 0.
  - (C) More than one solution if  $b = \exp(1)$ .
  - (D) A unique solution if  $a \neq 0$  and b > 0.
  - (E) None of the above.



- +=. The second-order linear differential equation  $(1-x^2)y''(x)+2xy'(x)-2y(x)=f(x)$ , for -1 < x < 1. To find the homogeneous solution, i.e. f(x)=0, given one solution  $y_1(x)=x$ , the other linearly independent solution  $y_2(x)$  can be derived by setting  $y_2(x)=v(x)y_1(x)$ . Assuming v(x) satisfies v(1)=2 and  $v(2)=\frac{5}{2}$ , which of the following statements about v(x) is/are true?
  - (A)  $x(1-x^2)v''(x) + 2v'(x) = 0$ .
  - (B)  $x(x^2 1)v''(x) + 2v'(x) = 0$ .
  - (C)  $v'(x) = \frac{x^2 1}{x^2}$ .
  - (D)  $v'(x) = \frac{x^2}{x^2-1}$ .
  - (E) None of the above.

類組:電機類 科目:工程數學 C(3005)

共12頁第8頁

※請在答案卡內作答

প্ত Continued from Problem  $+\Xi$ , find a particular solution for  $f(x)=1-x^2$  by the method of variation of parameters, i.e.,  $y_p(x)=u_1(x)y_1(x)+u_2(x)y_2(x)$ , where  $y_1(x)$  and  $y_2(x)$  are obtained from Problem  $+\Xi$ . Which of the following statements regarding  $u_1(x)$  and  $u_2(x)$  can be true?

(A) 
$$u_1(x) = \ln(1+x) - \ln(1-x) - x$$
.

(B) 
$$u_2(x) = \ln(1+x) + \ln(1-x)$$
.

(C) 
$$u_1(x) = x + \frac{x^3}{3}$$
.

(D) 
$$u_2(x) = -\frac{x^2}{2}$$
.



- +£. Solve the initial value problem of  $(2x-x^2)y''(x)-5(x-1)y'(x)-3y(x)=0$  with y(1)=0 and y'(1)=1 by power series of the form  $y(x)=\sum_{n=0}^{\infty}c_n(x-1)^n$ . Which of the following statements regarding the recurrence relation as well as values of coefficients  $c_n$  is/are true?
  - (A)  $c_{n+2} = \frac{n+2}{n+3}c_n$ .
  - (B)  $c_8 = 0$ .
  - (C)  $c_5 = \frac{8}{5}$ .
  - (D)  $c_9 = \frac{63}{128}$ .
  - (E) None of the above.

類組:電機類 科目:工程數學 C(3005)

共工及第一頁

※請在答案卡內作答

 $+\dot{\pi}$ . Let  $\underline{y}(x) = [y_1(x) \ y_2(x)]^{\top}$  and consider the following system of first-order differential equations

$$\underline{y}'(x) = \left[ egin{array}{cc} 6 & -7 \ 1 & -2 \end{array} 
ight] \underline{y}(x) + \left[ egin{array}{c} 2 \ 3 \end{array} 
ight]$$


Assuming  $\underline{y}(0) = [1 \ -1]^{\top}$ , let  $\underline{Y}(s) = [Y_1(s) \ Y_2(s)]^{\top} = \mathcal{L}\{\underline{y}(x)\}$ . Which of the following statements is/are true?

- (A)  $Y_1(1) = \frac{5}{8}$ .
- (B)  $Y_1(6) = \frac{85}{42}$ .
- (C)  $Y_1(7) Y_2(7) = \frac{11}{14}$ .
- (D)  $\frac{Y_2(8)}{Y_1(8)} = 0$ .
- (E) None of the above.



+ Continued from Problem +  $\stackrel{.}{\Rightarrow}$ , which of the following statements regarding the solution y(x) is/are true?

- (A)  $y_2'(0) = 6$ .
- (B)  $y_1''(0) = 8$ .
- (C)  $y_2''(0) = 3$ .
- (D)  $\lim_{x\to -\infty} (y_1(x) y_2(x)) = \frac{1}{4}$
- (E) None of the above.



類組:<u>電機類</u> 科目:<u>工程數學 C(3005)</u>

共<u>(</u> <sup>2</sup> 頁 第 <u>L D</u> 頁

※請在答案卡內作答

 $+ \wedge$  Let y(x) be a real-valued function satisfying the following second-order differential equation

$$y''(x) + y(x) = f(x)$$

Assume y(0) = y'(0) = 0 and  $f(x) = \sum_{n \ge 0} u(x - n\pi) \sin(x - n\pi)$ . Which of the following statements is/are true?

- (A) y(x) is a periodic function with period  $\pi$ .
- (B) y(x) has a Fourier-series representation for all x > 0.
- (C)  $y\left(\frac{\pi}{2}\right) = \frac{1}{2}$ .
- (D)  $y'\left(\frac{\pi}{2}\right) = \frac{\pi}{4}$ .
- (E) None of the above.



類組:電機類 科目:工程數學 C(3005)

共して頁第11頁

※請在答案卡內作答

+ h. Let  $f(x) = \frac{\pi - x}{2}$  and

$$g(x) = [f(x)(u(x) - u(x - \pi))] \star \left(\sum_{n = -\infty}^{\infty} \delta(x - n2\pi)\right)$$

It is known that g(x) has the following Fourier series representation

$$\tilde{g}(x) = \sum_{n=0}^{\infty} \left( a_n \cos \left( \frac{n2\pi}{T} x \right) + b_n \sin \left( \frac{n2\pi}{T} x \right) \right).$$

with minimal period T and Fourier series coefficients  $a_n$  and  $b_n$ . Which of the following statements is/are true?

- (A)  $a_4 = 0$ .
- (B)  $b_4 = \frac{1}{8}$ .
- (C)  $\tilde{g}(2\pi) = \frac{\pi}{4}$ .
- (D)  $\sum_{n=1}^{\infty} a_n^2 + b_n^2 = \frac{5}{96} \pi^2$ .
- (E) None of the above.



# 類組:電機類 科目:工程數學 C(3005)

# 共 12頁 第 12頁

# ※請在答案卡內作答

二十、 Continued from Problem 十九. Consider the following boundary value problem for the bivariate function y(x,t)

$$\frac{\partial}{\partial t}y(x,t) = \frac{\partial^2}{\partial x^2}y(x,t),$$

for  $x \in (0,\pi)$  with initial conditions  $y(0,t) = y(\pi,t) = 0$  and y(x,0) = xf(x), where f(x) is given in Problem  $+\pi$ . The solution y(x,t) can be representation in the following form

$$y(x,t) = \sum_{n \ge 0} c_n e^{-d_n t} \sin\left(e_n x\right)$$

for some  $c_n, d_n, e_n \in \mathbb{R}$ . Which of the following statements is/are true?

- (A)  $c_1 = \frac{4}{\pi}$ .
- (B)  $c_2 = 1$ .
- (C)  $d_3 = 3$ .
- (D)  $e_3 = 3$ .
- (E) None of the above.

